

In the claims:

Claims 1-25 (Cancelled).

26. (Currently amended) A method of preventing or treating a tumor in an animal which comprises administering to said animal a tumor inhibiting effective amount of an antigen-presenting cell, wherein said antigen presenting cell expresses at least one class I MHC or class II MHC determinant that is syngeneic to said animal and at least one class I or II MHC determinant that is allogeneic to said animal, and wherein said antigen-presenting cell is transformed transfected with genomic DNA isolated from the tumor cells of said animal.

Claims 27-40 (cancelled).

(Previously amended) The method according to claim 26, wherein said antigen presenting cell is further transformed transfected with a nucleic acid molecule coding for at least one cytokine.

A2. (Previously amended) The method of claim 41, wherein said cytokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, interleukin-7, interleukin-8, interleukin-9, interleukin-10, interleukin-11, interleukin-12, interferon-α, interferon-γ, tumor necrosis factor, granulocyte macrophage colony stimulating factor, and granulocyte colony stimulating factor.

Question (Previously amended) The method according to claim 26, wherein said antigenpresenting cell is selected from the group consisting of a fibroblast, a macrophage, a B cell, and a dendritic cell.

(Previously amended) The method according to claim 26, wherein said tumor is a solid tumor or a hematological tumor.

(Previously amended) The method of claim 44, wherein said tumor is selected from the group consisting of melanoma, lymphoma, plasmocytoma plasmacytoma, sarcoma, glioma, thymoma, leukemias, breast cancer, prostate cancer, colon cancer, esophageal cancer, brain cancer, lung cancer, ovary ovarian cancer, cervical cancer and hepatoma.

- **Q** 46. (Previously amended) The method according to claim 26, wherein said animal is a human subject.
- (New) A method of preventing a tumor recurrence in an animal which comprises 47. administering to said animal a tumor-inhibiting effective amount of an antigen-presenting cell, wherein said antigen presenting cell expresses at least one class I MHC or class II MHC determinant that is syngeneic to said animal and at least one class I or II MHC determinant that is allogeneic to said animal, and wherein said antigen-presenting cell is transfected with genomic DNA isolated from the tumor cells of said animal.
- (New) The method of claim 47, wherein said antigen presenting cell is further 48. transfected with a nucleic acid molecule coding for at least one cytokine.
- 49. (New) The method of claim 47, wherein said cytokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, interleukin-7, interleukin-8, interleukin-9, interleukin-10, interleukin-11, interleukin-12, interferon-α, interferon-γ, tumor necrosis factor, granulocyte macrophage colony stimulating factor, and granulocyte colony stimulating factor.
- 50. (New) The method of claim 47, wherein said antigen-presenting cell is selected from the group consisting of a fibroblast, a macrophage, a B cell, and a dendritic cell.